



Click Here for Thawing Instructions for semen frozen by SBS Labs

Help Line (877) 658-3328

Select Breeders Services

www.selectbreeders.com

copyright 2005 Select Breeders Service, Inc.

Management of Mares for Frozen Semen Insemination

For related information, video guides and research results visit www.selectbreeders.com

Prior to Insemination

- Verify that the mare is a suitable candidate for AI with frozen semen by performing a routine reproductive exam that may include culture, cytology and biopsy when indicated. Data suggests that aged (> 15 years) or repeatedly barren mares may have a reduced pregnancy rate and are not the best candidates for AI with frozen semen.
- Once the mare comes into estrus, palpate and/or scan daily to monitor follicular activity.
- Upon detection of a large (35-40 mm) pre-ovulatory follicle, administer hCG or deslorelin.

If only one dose of semen is available for insemination.

1. Examine the mare via ultrasound at 6 hour intervals starting 12-24 hours after hCG or deslorelin administration.
2. Inseminate the single dose of frozen semen as soon as ovulation is detected.

It is extremely important that mares being inseminated post-ovulation are inseminated within 6 hours of ovulation. A significant reduction in fertility will occur if mares are inseminated more than 6 hours post ovulation.

If more than one dose of semen is available during a given heat cycle.

1. Continue to examine the mare via ultrasound once daily and inseminate a single dose of frozen semen approximately 24 hours after hCG or deslorelin administration.
2. Examine the mare approximately 16 hours after insemination and inseminate a second dose of frozen semen even if the mare has already ovulated.
3. Examine the mare the following day to confirm ovulation. Insemination of a third dose may be required if the mare has still not ovulated.

A general goal for mares inseminated with frozen semen is to inseminate within 12 hours prior to and/or within 6 hours after ovulation. This protocol ensures that viable sperm are in the oviduct during that interval for any mare ovulating within a period of 18 to 52 hours following administration of hCG or deslorelin.

Suggested schedule for insemination of mares when more than one dose is available:

Daily examinations during estrus (any time)

Day 0 Day of 35 to 40 mm follicle detection

Administer hCG or deslorelin at approximately 4:00 PM

Day 1 Inseminate a single dose of frozen semen at 4:00 PM (24 hrs post-injection).

Day 2 Inseminate a second dose of frozen semen at 8:00 AM (40 hours post-injection).

Day 3 Examine to confirm ovulation & inseminate a third dose if the mare has not ovulated at 8:00 AM

